No.



8400027

TO) ARE TO) WHOM: THESE: PRESENTS: SHARE, COME:

Kansas Agricultural Experiment Station

TUltereas, there has been presented to the

Secretary of Agriculture

AN APPLICATION REQUESTING A CERTIFICATE OF PROTECTION FOR AN ALLEGED NOVEL VARIETY OF SEXUALLY REPRODUCED PLANT, THE NAME AND DESCRIPTION OF WHICH ARE CONTAINED IN THE APPLICATION AND EXHIBITS, A COPY OF WHICH IS HEREUNTO ANNEXED AND MADE A PART HEREOF, AND THE VARIOUS REQUIREMENTS OF LAW IN SUCH CASES MADE AND PROVIDED HAVE BEEN COMPLIED WITH, AND THE TITLE THERETO IS, FROM THE RECORDS OF THE PLANT VARIETY PROTECTION OFFICE, IN THE APPLICANT(S) INDICATED IN THE SAID COPY, AND WHEREAS, UPON DUE EXAMINATION MADE, THE SAID APPLICANT(S) IS (ARE) ADJUDGED TO BE ENTITLED TO A CERTIFICATE OF PLANT VARIETY PROTECTION UNDER THE LAW.

NOW, THEREFORE, THIS CERTIFICATE OF PLANT VARIETY PROTECTION IS TO GRANT UNTO THE SAID APPLICANT(S) AND THE SUCCESSORS, HEIRS OR ASSIGNS OF THE SAID APPLI-CANT(S) FOR THE TERM OF eighteen* YEARS FROM THE DATE OF THIS GRANT, SUBJECT TO THE PAYMENT OF THE REQUIRED FEES AND PERIODIC REPLENISHMENT OF VIABLE BASIC ED OF THE VARIETY IN A PUBLIC REPOSITORY AS PROVIDED BY LAW, THE RIGHT TO EX-OTHERS FROM SELLING THE VARIETY, OR OFFERING IT FOR SALE, OR REPRODUCING IT. RTING IT, OR EXPORTING IT, OR USING IT IN PRODUCING A HYBRID OR DIFFERENT HEREFROM, TO THE EXTENT PROVIDED BY THE PLANT VARIETY PROTECTION ACT. TED STATES SEED OF THIS VARIETY (1) SHALL BE SOLD BY VARIETY NAME ONLY AS CERTIFIED SEED AND (2) SHALL CONFORM TO THE NUMBER OF GENERATIONS HE OWNER OF THE RIGHTS. (84 STAT. 1542, AS AMENDED, 7 U.S.C. 2321 ET SEQ.)

AMENDED CERTIFICATE
*Original grant January 31, 1986.
In Lestimony Winercot, I have hexeunto set my hand and caused the seal of the Blaut Variety Protection Office to be affixed at the City of Washington, D. C.

the year of our Lord one thousand nine hundred and ninety-one.

Plant Variety Protection Office ricultural Marketing Service

U.S. DEPARTMENT OF AGRICULTURE AGRICULTURAL MARKETING SERVICE LIVESTOCK, MEAT, GRAIN & SEED DIVISION			FORM APPROVED: OMB NO.0581-0055 No certificate for plant variety protection		
(Instruction	ns on reverse)		553).		
1. NAME OF APPLICANTIS) KAUSAS AGRICU		2. TEMPORARY DESIGNATION	3. VARIETY NAME		
Pioneer Hi-Bred International, Plant Breeding Division EXPER Dept. of Cereal Seed Breeding	INGNT STATE	W7442B	2157		
4. ADDRESS (Street and No. or R.F.D. No., City, Sta	ate, and Zip Code)	5. PHONE (Include area code)	FOR OFFICIAL USE ONLY		
WATERS HALL KANSAS STATE UNIL	ERSITY	(913) 532-6147	8400027		
Rt. 2 MANHATTAN, KS 66506-4008 Hutchinson, Kansas 67501		(316) 662 5439	0400027		
6. GENUS AND SPECIES NAME	7. FAMILY NA	ME (Botanical)	DATE		
			2 12/23/83		
Triticum aestivum	gramine		= TIME 2:30 □ A.M.		
	1 <u> </u>		AMOUNT FOR FILING		
8. KIND NAME]9.	DATE OF DETERMINATION	1 000		
Wheat		September 1, 1981	DATE		
WIIEGL		pehremmer I, 130T	DATE 12/23/83 AMOUNT FOR CERTIFICATE		
10. If THE APPLICANT NAMED IS NOT A "PERSO partnership, association, etc.)					
partiters inp, association, etc.)			\$ 500.00 H DATE		
Corporation			12/27/85		
11. IF INCORPORATED, GIVE STATE OF INCORP	ORATION		12. DATE OF INCORPORATION		
Iowa		•	May, 1926		
13. NAME AND ADDRESS OF APPLICANT REPRE	SENATIVE(S), II ハケル	FANY, TO SERVE IN THIS APPLICA NON A. SCHAFFER	ATION AND RECEIVE ALL PAPERS		
Dr. Charles Hayward Pioneer Hi-Bred International,	20-25	ARTMENT OF AGRONOM)	?		
Rt. 2	THRO	CKMORTON HALL	and the second of the second o		
Hutchinson, Kansas 67501		SAS STATE UNIVERSETY HATTAN, KS 66506-550			
14. CHECK APPROPRIATE BOX FOR EACH ATTA					
a. X Exhibit A, Origin and Breeding History of the		c. Exhibit C, Objective D	escription of the Variety (Request form		
Section 52 of the Plant Variety Protection A	(21.)	in from Films Variety Fro	decision officery		
b. X Exhibit B, Novelty Statement		d. X Exhibit D, Additional	Description of the Variety		
15. DOES THE APPLICANT(S) SPECIFY THAT SEE	D OF THIS VAR	IETY BE SOLD BY VARIETY NAME	ONLY AS A CLASS OF CERTIFIED		
SEED? (See Section 83(a) of the Plant Variety Pi	otection Act.).		tems 16 and 17 below) No		
16. DOES THE APPLICANT(S) SPECIFY THAT THI LIMITED AS TO NUMBER OF GENERATIONS:		17. IF "YES" TO ITEM 16, V BEYOND BREEDER SEE	WHICH CLASSES OF PRODUCTION D?		
Yes No		Foundation	Registered Certified		
18. DID THE APPLICANT(S) FILE FOR PROTECT	ON OF THE VAF	RIETY IN THE U.S. OR OTHER COU	Yes (If "Yes," give name		
		•	of countries and dates)		
			X No		
19. HAVE RIGHTS BEEN GRANTED IN THE U.S. (OR OTHER COUP	NTRIES?	Yes (If "Yes," give name		
		-	of countries and dates)		
	•		No No		
20. The applicant(s) declare(s) that a viable sam	ple of basic seed	ds of this variety will be furnished	with the application and will be re-		
plenished upon request in accordance with s			iety and helieveled that the variety is		
The undersigned applicant(s) is (are) the ow distinct, uniform, and stable as required in S Variety Protection Act.	Section 41, and	is entitled to protection under the	e provisions of Section 42 of the Plant		
Applicant(s) is (are) informed that false repr	esentation here	in can jeopardize protection and i	result in penalties.		
SIGNATURE OF APPLICANT	DATE				
Pioneer Hi-Bred International, I					
by Charles To Hayword	December 8, 1983				
SIGNATURE OF APPLICANT			DATE		

States explain

18/4/2/

FORM LMGS-470 (9-81) (Edition of 1-78 is obsolete)

2157, Triticum aestivum L. em Thell, a hard red winter wheat, was developed by Pioneer Hi-Bred International, Inc. from the cross of Caprock/4/Etoile de Choisy//Thorne/Clarkan/3/CI13390/5/SC3212. A semidwarf mutant was selected from the Etoile de Choisy//Thorne/Clarkan cross. This mutant was crossed to CI13390. A pure line selection from this cross was crossed to Caprock. This single cross was then crossed to SC3212. SC3212 is a pure line resulting from a Sturdy outcross.

The F_1 generation was grown in the greenhouse at Hutchinson in 1971. Single F_2 plants were selected in 1973 and the bulked seed of each plant was advanced in the field at Hutchinson in 1974. Heads were selected from the 1974 bulk and planted as individual rows in 1975. Seed from a single F_4 -derived row was bulked and was advanced from the F_5 to the F_7 at Hutchinson. Single heads were selected from the F_7 bulk in 1978. The F_7 -derived seed from one head was grown in a row in 1979 and the bulked progeny plot from this row was selected in 1980 near Hastings, Nebraska. 2157 has been in yield tests and milling and baking trials since 1981.

2157 has shown uniformity and stability for all traits as described in Exhibit C (Form LPGS-470-6) -- "Objective Description of Variety."

No variants have been observed in 2157.

14B. Exhibit B, Novelty Statement

2157 is an awned semidwarf hard red winter wheat cultivar, most similar to the variety Sturdy in many phenotypic and agronomic traits. 2157 is uniquely different from Sturdy in leaf carriage and in Hessian fly resistance. 2157 has an erect flag leaf at booting, whereas Sturdy has a recurved flag leaf. Also, Sturdy is susceptible to all races of Hessian fly while 2157 is resistant to races GP, A, B, and C.

2157 averaged about 99 cm in height at Hutchinson, Kansas, in 1983, about 2 cm taller than Sturdy but 21 cm shorter than Scout. In Pioneer trials, 2157 has been more winterhardy than Sturdy.

2157 and Sturdy show adult plant resistance to current races of leaf rust (*Puccinia recondita* Rob. ex Desm. f. sp. tritici) but little resistance to current races of stem rust (*P. graminis* Pers. f. sp. tritici Eriks) in Kansas.

EXHIBIT C (Wheat)

U.S. DEPARTMENT OF AGRICULTURE AGRICULTURAL MARKETING SERVICE LIVESTOCK, POULTRY, GRAIN & SEED DIVISION BELTSVILLE, MARYLAND 20705

OBJECTIVE DESCRIPTION OF VARIETY

INSTRUCTIONS: See Reverse. WHEAT (TRITICUM SPP.)	
NAME OF APPLICANTIS	FOR OFFICIAL USE ONLY
Pioneer Hi-Bred International, Inc.	PVPO NUMBER
ADDRESS (Street and No. or R.F.D. No., City, State, and ZIP Code) Plant Breeding Division Department of Cereal Seed Breeding	8400027 VARIETY NAME OR TEMPORARY DESIGNATION
Rt. 2 Hutchinson, Kansas 67501	
Place the appropriate number that describes the varietal character of this variety in the Place a zero in first box (e.s. 0 8 9 or 0 9) when number is either 29 or less or	boxes below.
1. KIND:	A Secretary of the Control of the Co
1 = COMMON 2 = DURUM 3 = EMMER 4 = SPELT 5 = POLISH 6 = POUI	LARD 7 = CLUB
2. TYPE:	
2 1 = SPRING 2 = WINTER 3 = OTHER (Specify) 2 = HARD	3 = OTHER (Specify)
2] = WHITE 2 = RED 3 = OTHER (Specify)	The Statement of the Control of the
3. SEASON - NUMBER OF DAYS FROM EMERGENCE TO:	· · · · · · · · · · · · · · · · · · ·
2 3 0 FIRST FLOWERING 2 3 7 LAST	FLOWERING
4. MATURITY (50% Flowering):	
0 5 NO. OF DAYS EARLIER THAN	2 = SCOUT 3 = CHRIS
NO. OF DAYS LATER THAN	5 = NUGAINES 4.76 = LEEDS
5. PLANT HEIGHT (From soil level to top of head):	
0 9 9 CM. HIGH	and the second of the second o
CM. TALLER THAN	3 = CHRIS
2 1 CM. SHORTER THAN	2 = SCOUT 3 = CHRIS 5 = NUGAINES 6 = LEEDS
S. PLANT COLOR AT BOOTING (See reverse): 7. ANTHER COLOR:	
3 1 = YELLOW GREEN 2 = GREEN 3 = BLUE GREEN 1 1 = YELLOW	2 = PURPLE
3. STEM:	
1 Anthocyanin: 1 = ABSENT 2 = PRESENT 2 Waxy bloom: 1 :	ABSENT 2 = PRESENT
Hairiness of last internode of rachis: 1 = ABSENT 2 = PRESENT 1 Internodes: 1 =	HOLLOW 2 = SOLID
0 4 NO. OF NODES (Originating from node above ground) 2 7 CM. INTER	NODE LENGTH BETWEEN FLAG LEAF BELOW
. AURICLES:	
1 Anthocyanins 1 = ABSENT 2 = PRESENT 1 Hairiness: 1 = A	ABSENT 2 = PRESENT
D. LEAF:	
Flag leaf at = ERECT 2 = RECURVED 1 Flag leaf: = N	OT TWISTED 2 = TWISTED
1 Hairs of first leaf sheath: 1 = ABSENT 2 = PRESENT 2 Waxy bloom of fli	ag leaf sheath: 1 = ABSENT 2 = PRESENT
0 9 MM. LEAF WIDTH (First loaf below flag load) 2 2 CM. LEAF	LENGTH (First leaf below flag leaf);
DBM LPGS.470.6 (3.79) (Formerly Form GR.470.6 (2.73) which may be used)	

	<u> </u>	the state of the s
11. HEAD: 1 Density: 1 = LAX	2 - DENSE	Shape: 1 = TAPERING 2 = STRAP 3 = CLAVATE 4 = OTHER (Specify) Fusiform
4 Awnedness: 1 = AWNL	ESS 2 = APICALLY AWNLETED 3 =	The state of the s
Color at maturity: 5 =	WHITE 2 YELLOW 3 PINK 4 =	
0 9 CM. LENGTH	The second of th	0 9 мм. wіртн
12. GLUMES AT MATURITY Length: 1 = SHORT (C. 3 = LONG (C.	Ys, 2A, 7 mm.) 2 = MEDIUM (CA, 8 mm.) A.(9 mm.)	Width: 1 = NARROW (CA. 3 mm.) 2 = MEDIUM (CA. 3.5 mm.) 3 = WIDE (CA. 4 mm.)
Shoulder I = WANTIN shape: 4 = SQUARE	G 2 = OBLIQUE 3 = ROUNDED 5 = ELEVATED 6 = APICULATE	3 Beak: 1 = OBTUSE 2 = ACUTE 3 = ACUMINATE
13. COLEOPTILE COLOR:		14. SEEDLING ANTHOCYANIN: 1
· · · · · · · · · · · · · · · · · · ·	3 = PURPLE	I I - ABSENT 2 - FRESENT
15. JUVENILE PLANT GRO	WTH HABIT:	
2 1 = PROSTRATE.	2 = SEMI-ERECT 3 = ERECT	
16. SEED:		
3 Shape: To OVATE	2 = OVAL - 3 ELLIPTICAL	1 Cheek: N= ROUNDED 2 = ANGULAR
2 Brush, 1 = SHORT	2 - MEDIUM 3 : LONG	1 Brush: 1 = NOT COLLARED 2 2 = COLLARED
Phenol reaction	\$ \$ = (Vory > 2 = FAWN > 3 = LT. BROWN	
3 (See instructions):	4 = BROWN 5 = BLACK	
3 Color: 1 = WHITE	2 = AMBER 3 = RED 4 = PURPLE	5 = OTHER (Specify)
0 6 MM. LENGTH	0 3 мм. WIDTH	2 8 GM. PER 1000 SEEDS
17. SEED CREASE: Width: = 60% OR LE	ESS OF KERNEL 'WINOKA'	Depth: 1 = 20% OR LESS OF KERNEL 'SCOUT'
	SS OF KERNEL 'CHRIS'	1 2 = 35% OR LESS OF KERNEL 'CHRIS'
3 = NEARLY A	WIDE AS KERNEL 'LEMHI'	3 = 50% OR LESS OF KERNEL 'LEMHI'
	ed, 1 = Susceptible, 2 = Resistant)	
O STEM RUST (Races)	LEAF RUST (Races)	O STRIPE RUST O LOOSE SMUT
1 POWDERY MILDEW	0 BUNT	OTHER (Specify)
19. INSECT: (0 = Not Teste	d, 1 = Susceptible, 2 = Resistant)	
0 SAWFLY	1 APHID (Bydy.)	O GREEN BUG O CEREAL LEAF BEETLE
OTHER (Specify)	HESSIAN FLY	2 GP 2 A 2 2 B 2 C
	RACES	
ting the second		0 D 0 E 0 F 0 G
	TY MOST CLOSELY RESEMBLES THAT S	UBMITTED:
CHARACTER	NAME OF VARIETY	CHARACTER NAME OF VARIETY
Plant tillering	Sturdy	Seed size Vona
Leof size	Sturdy	Seed shape Sturdy
Leaf color	Tam W-101	Coleoptile elongation Eagle
Leaf carriage	Tam 105	Seedling pigmentation PL145

INSTRUCTIONS

GENERAL: The following publications may be used as a reference aid for the standardization of terms and procedures for completing this form:

- (a) L.W. Briggle and L. P. Reitz, 1963, Classification of Triticum Species and Wheat Varieties Grown in the United States, Technical Bulletin 1278, United States Department of Agriculture.
- (b) W.E. Walls, 1965, A Standardized Phenol Method for Testing Wheat Seeds for Varietal Purity, contribution No. 28 to the handbook of seed testing prepared by the Association of Official Seed Analysts. (See attachment.)

LEAF COLOR: Nickerson's or any recognized color fan should be used to determine the leaf color of the described variety.

14D. Exhibit D, Additional Description of the Variety

2157 is a common hard red winter wheat, Triticum aestivum L.

Flowering date of 2157 is 5 days earlier than the variety Scout and 3 to 4 days earlier than Newton. At Hutchinson, Kansas, when seeded about September 30, average first flowering is about 230 days after emergence. Last flowering averages about 7 days later. It is recognized that environmental factors influence flowering of varieties differently.

2157 has averaged 99 cm in height, about 8 cm shorter than Newton and 21 cm shorter than Scout.

The plant color of 2157 at booting stage is blue green.

Anther color of 2157 is yellow, similar to Scout and Newton.

Anthocyanin has been absent in the stem of 2157. There is a waxy bloom on the stem. Internodes of 2157 are hollow. At maturity, stems are yellow and very strong. Normally 4 stem nodes are present above ground. Internode length between flag leaf and leaf below is about 27 cm.

Auricles of 2157 are glabrous and lacking in anthocyanin.

Flag leaves are generally erect at booting and tend not to be twisted. Hairs have not been observed on the first leaf sheath. A moderate amount of waxy bloom occurs on the last leaf sheath. The first leaf below the flag leaf averages about 9 mm wide and 22 cm long when measured at Hutchinson, Kansas, in 1983.

Spikes are generally mid-dense to lax, fusiform, awned, and white at maturity. Awns are rough and about 5 to 7 cm in length. Spike width and length averages about 9 mm and 9 cm, respectively. However, spike width and length are variable with plant population and level of production.

The glumes of 2157 are short and narrow with generally square shoulders, and glabrous. Beaks are acuminate and range from 2 to 4 mm long.

When evaluated at Hutchinson, Kansas, coleoptile color is white and seedling anthocyanin is absent.

Kernels are red in color, elliptical to ovate in shape, with rounded cheeks and a shallow crease. The brush is not collared and medium in size. The embryo is medium in size. Kernels average 6 mm long and 3 mm wide and about 28 g per 1000. Phenol reaction is light brown.

2157 is moderately resistant to soil borne mosaic virus. It is moderately resistant to leaf rust (Puccinia recondita Rob. ex Desm. f. sp. tritici) and moderately susceptible to stem rust (P. graminis Pers. f. sp. tritici Eriks) races currently common in Kansas. 2157 has not been tested for specific races of leaf rust. 2157 is resistant to GP, A, B, and C of Hessian fly (Mayetiola destructor Say). It is susceptible to powdery mildew (Erysiphe graminis f. sp. tritici E. Marchal). 2157 has not been tested for bunt (Tilletia caries (DC.) Tul.) resistance.

2157 has a good yield record when compared with currently grown hard red winter wheats (Table 1). In the presence of *Septoria* spp., 2157 has a pronounced yield advantage over susceptible varieties. Short plant height and superior straw strength give 2157 excellent resistance to lodging.

2157 has good quality characteristics when compared to the average of current standard varieties in the HRW area. It has been tested by the Pioneer Wheat Quality Laboratory for several years. Table 2 gives 1981-83 values for 2157, Newton, Centurk 78, Vona, and all checks grown in plots tested in the Great Plains. Each value is an average of eight tests. The figures show that 2157 is superior to the checks for flour yield and protein while it is slightly poorer for mixing time and tolerance as evaluated with the mixograph. Loaf volume is comparable to Centurk 78 and Vona, but lower than the average of all checks.

Table 1. Yield comparisons of 2157 with standard varieties tested in 1981-82 and 1982-83. Data represents average of 104 replications.

Variety	Yield (bu/acre)
2157	56
Centurk 78	50
Newton	52
Tam 105	54
Triumph 64	47
Vona	. 53

Table 2. Milling and baking evaluation of 2157.

Variety	% Flour Yield	% Flour Protein	Mixo Mx Time	ograph Tolerance	Microbake (10 g) Loaf Volume (cc)
2157	69.3	11.4	3.4	2.9	57.5
Centurk 78	67.7	10.5	3.8	6.4	57.0
Newton	66.4	10.7	3.2	3.9	62.5
Vona	67.3	10.1	3.6	5.4	58.5
All checks	67.9	10.8	3.4	4.6	60.0

 $^{^1}$ Mixograph mixing tolerance--evaluated on a 9-point scale on which 9 = very good, 5 = fair, 1 = very poor

14E. Exhibit E, Statement of the Basis of Applicant's Ownership

Pioneer Hi-Bred International, Inc., Plant Breeding Division, believes it is the sole, original and first breeder of the 2157 variety of hard red winter wheat for which it solicits a certification of protection.